



11TH EUROPEAN CONGRESS ON TROPICAL MEDICINE AND INTERNATIONAL HEALTH

16-20 SEPTEMBER 2019
LIVERPOOL, UK

The clinical sensitivity of standard direct wet mount microscopy for soil-transmitted helminth infections in school children in Jimma, South-West Ethiopia

Daniel Dana^{1,3}, Johnny Vlamincx², Zeleke Mekonnen¹, Mio Ayana¹, Florian Vogt³, Kristien Verdonck³, Hannock Tweya⁴, Collins Timire⁴, Peter Geldhof², Bruno Levecke²

¹School of Medical Laboratory Science, Faculty of Health Science, Institute of Health, Jimma University, Jimma, Ethiopia

²Department of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium

³Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

⁴The International Union against Tuberculosis and Lung Disease, Paris, France

Abstract

Introduction: Soil-transmitted helminthiasis is a disease caused by *Ascaris lumbricoides*, *Trichuris trichiura* and the two hookworms (*Ancylostoma duodenale* and *Necator americanus*) and remains a major public health problem in school children in Ethiopia. Although direct wet mount microscopy is the standard means to diagnose intestinal parasitic diseases in health care facilities in Ethiopia, it remains unclear what its diagnostic performance is for soil-transmitted helminths.

Aim: To determine the clinical sensitivity of standard direct wet mount microscopy for soil-transmitted helminth infections compared to the composite reference standard in school children in Jimma, South-West Ethiopia

Methods: A school based cross-sectional study was performed in Jimma Town and included 600 children from 10 primary schools. The clinical sensitivity of direct wet mount microscopy was compared to a composite reference standard consisting of Kato-Katz thick smear, McMaster and Mini-FLOTAC.

Results: The composite reference standard about 210 *Ascaris* (35.0%), 312 *Trichuris* (52.0%) and 102 hookworm cases (17.0%). The median intensity of infections (expressed as eggs per gram of stool (EPG)) equalled 2,057 EPG for *Ascaris*, 200 EPG for *Trichuris* and 110 EPG for hookworms. The sensitivity of direct wet mount microscopy was 73.8% for *Ascaris*, but was only around 17% for both *Trichuris* and hookworms. For each of the three soil transmitted helminths there was an increase in sensitivity as a function of increasing intensity of infections. For *Ascaris*, the sensitivity ranged from 30.2% for the lowest infection intensities to 94.3% for the highest levels of infection intensity. For both *Trichuris* and hookworms, the sensitivity ranged from nearly 0% to 42.9% across the different levels of infection intensity.

Conclusion: Using direct wet mount microscopy severely underestimated the prevalence of soil transmitted helminths. The sensitivity of direct wet mount microscopy is low for STH, particularly for infections with *Trichuris* and hookworm when the infection intensities are low.